

Severn Pathology Infection Sciences Sample Quick Guide

This quick reference guide for commonly sent samples is part of Infection Sciences' commitment to improve patient care and environmental sustainability by attempting to reduce rejected specimens. Many specimens are rejected for testing due to the use of an incorrect specimen container. This could result in a delay in diagnosis and treatment for patients and leads to unnecessary waste and transport costs.

Please ensure all samples:

- Are correctly identified with NHS number/hospital number, DoB, name and site of sampling or specimen type.
- Contain correct clinical details including **FOREIGN TRAVEL**, antimicrobial therapy, sample date, and duration of illness (important for serology).
- Have their hazard risk included on all requests (**including foreign travel, contact with imported animals and consumption of unpasteurised milk**) irrespective of test requested. **This information is essential to ensure that the correct laboratory precautions are undertaken when processing specimens. Please contact the Infection Sciences Laboratory prior to sending samples in patients at risk of high risk organisms. Please refer to the Infection Sciences Laboratory User manual for further information.**
- Go in the correct coloured bag - **Blue for Infection Sciences**. Please ensure samples are not leaking and the lid is secure.
- For electronically requested specimens, attach barcode labels over the tube label, with the **top of the barcode placed directly below the tube lid**.









To request testing kits and tubes please email
pathologyconsumablesouthmead@nbt.nhs.uk



or order from pathology stores at <https://www.nbt.nhs.uk/severn-pathology/requesting/consumables-ordering>





For any other queries or suggestions for improvements relating to the below please email microbiology@nbt.nhs.uk or ISQuality@nbt.nhs.uk

For a full list of Pathology testing repertoire including turnaround times please see [Test Information | North Bristol NHS Trust \(nbt.nhs.uk\)](#) (Last updated December 2024)

Blood cultures (Adults)	8-10mls blood in each bottle. 2 Sets of cultures at separate sites should be obtained (other than line or infective endocarditis)	
Blood culture (Paediatrics)	1-3mls blood	
Swab for bacterial/candida culture Ears/ eyes/ wounds/ nose/ throat/ high vaginal/ lower vaginal/ cervical/ self-taken (For Chlamydia, Gonorrhoea, Trichomonas, M. genitalium NAAT send an Aptima tube) (See below)	Purple top liquid swab	






<p>Chlamydia, Gonorrhoea, Bacterial Vaginosis (Trichomonas and M. genitalium by special request only) NAAT swab (Vaginal/ Eye)</p>	<p>Aptima (use Vaginal) swab. Please check expiry date of kit.</p>	
<p>Chlamydia and Gonorrhoea NAAT urine</p>	<p>Preferred sample for males. Please do not under/overfill SAMPLE SHOULD BE IN BETWEEN THE TWO BLACK LINES. Please check expiry date of tube.</p>	
<p>MRSA screen swab</p>	<p>Purple top sigma Transwab White top red/white sigma Duo Transwab If needing to screen urine, send boric acid container</p>	 <p>Order Details Σ-Transwab® MW176S NHS Supply Chain Code: HHD260</p> <p>HHD271 Σ-Transwab® Duo MW167S</p>


<p>Carbapenemase screen (CPE) / VRE screen (rectal swab)</p>	<p>MUST have faecal matter on the rectal swab otherwise cannot be process and will be rejected</p>	
<p>Viral Nose and Throat swabs/ Vesicles/ Eyes/ Genital lesions/ M pox</p>	<p>Red top Copan tube Green top universal transport medium (UTM) tube Purple top UTM tube White top UTM tube</p>	

<p>Bordetella pertussis</p> <p>Viral Nose and Throat swabs</p> <p>Nasopharyngeal swab for PCR</p> <p>Please see UKHSA for advice on testing Pertussis: guidelines for public health management - GOV.UK (www.gov.uk)</p>	<p>Red top Copan tube</p> <p>Green top universal transport medium (UTM) tube</p>	
<p>Stool</p> <p>Bacterial and viral enteric PCR / C. difficile PCR</p> <p>Viral gastroenteritis and Norovirus PCR</p> <p>Ova, cysts, & parasites (OCP)</p> <p>H pylori antigen</p>	<p>Blue topped with a scoop, minimum 1/3 full. Ideally separate samples for each ICE request sticker.</p> <p>Please postdate OCP samples with different days otherwise they may be rejected by the laboratory.</p>	
<p>Threadworm/ Enterobius kit</p>	<p>Rectal swab suspension</p>	
<p>CSF MC&S/ Biofire PCR/ CRAG antigen</p>	<p>Universal white top container</p>	

Measles	Oral fluid testing - please contact UKHSA to request a kit 0300 3038162	
Measles PCR	<p>If wanting to test in clinic, please discuss first with UKHSA to risk assess</p> <p>Red top Copan tube</p> <p>Green top universal transport medium (UTM) tube</p>	
<p>Mumps (oral fluid IgM /PCR)</p> <p>Acute cases only.</p> <p>For immunity testing please see viral serology below</p>	<p>Oral fluid testing - please contact UKHSA to request a kit 0300 3038162</p> <p>If wanting to test in clinic, please discuss with Virology at Severn Infection Sciences to avoid delay</p>	
Sputum MC&S	Universal white top container	
<p>Bronchoscopy for MC&S/</p> <p>Virology PCR/ PCP PCR/</p> <p>Aspergillus antigen</p>	Universal white top container	

Tissue for MC&S/ Peritoneal dialysis fluid/ Pus sample	Universal white top container	
Urine MC&S if >2ml	Boric Acid container Olive Green/Red	
Urine MC&S if < 2ml i.e. paediatric samples	White top universal container N.B. If sample is >2ml this will be rejected by the laboratory	
Urine for Legionella and Pneumococcal antigens	White top universal container	
Blood for bacterial/viral serology e.g. Lyme/ CMV/ EBV/ Hepatitis/ HIV/ Varicella Zoster/ Measles	Gold top tube Do not send a green top tube	

<p>Blood for viral PCR</p> <p>e.g. CMV/ EBV/ HIV Viral load/ Hepatitis B, C</p>	<p>Purple top EDTA tube</p>	
<p>Antimicrobial assays, except for Isoniazid (+ N-acetyl-Isoniazid) and Azoles</p>	<p>Gold top tube</p>	
<p>Isoniazid (+ N-acetyl-Isoniazid)</p>	<p>Grey top tube (fluoride oxalate; unspun whole blood)</p>	
<p>Azoles (itraconazole, posaconazole, voriconazole, isavuconazole, fluconazole)</p>	<p>Red top tube (plain clotted blood)</p> <p>Or Gold top tube[†]</p> <p>[†]If a gel tube is used, separate serum into a secondary tube as soon as possible as the gel may bind the analyte</p>	 

Blood for fungal markers, beta glucan/ Aspergillus antigen	Gold top tube	
Fungal skin/ nail scraping for microscopy and culture	Mycology collection kit	